CORRECTION AND SCOPE

Global ELF Fields and Biological Synchrony: An Exploratory Analysis

Exploratory interdisciplinary analysis. The reported correlations cannot establish the proposed quantum or whole-body mechanisms. Those mechanisms require independent evidence.

Manuscript · p. 1, introduction

Evidence · Live publication library

ChatGPT

Global ELF Fields and Biological Synchrony: An Exploratory Analysis

Introduction



Conjectured multi-level interactions between Schumann Resonances (SRs) and the human body. Brain and heart synchronization with SRs has been observed in research, and preliminary evidence hints at links to other organ systems. At quantum scales, photon-based mechanisms (lower red oval) are hypothesized as a likely means for SRs to influence biological processes [1] [2] .

Our planet is constantly enveloped by extremely low-frequency (ELF) electromagnetic fields. Chief among these are the *Schumann resonances* (SRs) – standing electromagnetic waves in the Earth-ionosphere cavity excited by global lightning activity. The fundamental Schumann mode occurs at about 7.8 Hz (with higher harmonics near 14, 20, 26 Hz, etc.), corresponding to an electromagnetic wavelength roughly equal to Earth's circumference [3] . Intriguingly, these frequencies overlap with the characteristic electrical oscillations of the mammalian brain. For example, the fundamental Schumann frequency (~7–8 Hz) lies in the theta–alpha range of brain waves, and was famously noted to coincide with human α-rhythm around 7.5–13 Hz [4] . This coincidence suggests a potential for resonance or entrainment between global ELF fields and neural activity, a hypothesis first raised in the 1950s and still provoking scientific curiosity.

The key question explored in this paper is: **Can global ELF/VLF electromagnetic fields (particularly Schumann resonances) weakly bias or coordinate biological behavior on large scales?** We assume **a priori** that these fields *do* interact with nervous systems (human and non-human), albeit weakly, and examine how such weak coupling might be amplified or integrated into significant biological effects. We will

discuss mechanisms like stochastic resonance (noise-assisted signal detection) and massive parallelism in neural ensembles, which could allow ultra-weak periodic influences to accumulate into measurable outcomes over long timescales and across large populations. We then outline a physically and biologically grounded model of ELF–brain interaction – involving threshold biasing of neural activity, neuronal phase entrainment, and slow ensemble drifts – and review empirical research spanning physics, neuroscience, and systems biology. Throughout, the tone remains scientifically exploratory and grounded in known principles, with proposed experiments and falsifiable predictions to guide future research.

Background: Schumann Resonances and Bioelectromagnetic Rhythms

What are Schumann resonances? Physically, SRs are global standing waves of ELF electromagnetic energy trapped between the Earth's surface and the conductive ionosphere about 100 km above [5] [6]. The Earth–ionosphere cavity behaves like a spherical resonator. Lightning strikes around the world excite this cavity, launching pulses that circumnavigate and interfere. At certain frequencies determined by the Earth's size and ionospheric properties, resonance conditions occur, producing stable modes [7]. The fundamental mode (~7.8 Hz) corresponds to a wavelength approximately equal to Earth's circumference (~40,000 km) [8] [3]. Higher harmonics near 14, 20, 26, 33 Hz and beyond are also observed, though with diminishing intensity [3]. These resonant frequencies remain nearly constant (changing only if Earth's cavity properties change) and provide a perpetual background "hum" of ELF electromagnetic oscillations worldwide. Importantly, ELF waves penetrate matter far better than higher frequencies; a 7–8 Hz EM wave can propagate through seawater to depths of order 100 m [9]. Thus, Schumann resonances have enveloped *all* living environments – land, ocean, and probably subsurface habitats – since life's emergence [10] [9].

Biological frequencies and evolutionary exposure: The fact that Schumann resonance frequencies overlap with neural rhythm frequencies (theta, alpha, beta, etc.) has long raised the idea that organisms might have *evolved* under this global EM "heartbeat" and possibly even utilize it. The German scientist W.O. Schumann and physician Herbert König first noted the match between 7.83 Hz and human brainwaves, suggesting a resonance between the "Earth's brain" and our own [4]. Subsequent research indeed found that the human brain's natural resonant frequency appears to be around 10 Hz (within the alpha band) and varies slightly with skull size [11]. This is intriguingly close to the first SR modes. Such observations have led to speculation that living systems might have *tuned in* to these ever-present signals as a timing reference or an informational cue over evolutionary timescales [10] [12]. In other words, global ELF fields could play a role analogous to light–dark cycles or other environmental rhythms in influencing biological clocks and behaviors. In fact, evidence exists that an **absence** of the Schumann-frequency field can disturb biological rhythms: in the 1960s–70s, Rütger Wever isolated human volunteers in an underground bunker shielded from natural EM fields and observed their circadian cycles desynchronize from 24 hours; remarkably, introducing a weak 10 Hz electromagnetic signal (approximate SR frequency) into the bunker helped re-synchronize the subjects' rhythms [13]. This classic experiment suggests that Schumann-range signals may act as a "zeitgeber" (time-giver) for the circadian system, or at least that the human body has some receptivity to ELF cues. Wever's work and related studies led some researchers to call the 7–8 Hz SR wave a "natural circadian tuning signal" for life [14] [15]. While the primary circadian entrainment cue is light, these results hint that the Earth's electromagnetic background might serve as a secondary environmental reference that organisms have historically leaned on for maintaining physiological harmony with the planet's cycles.

Field intensities and the challenge of weak coupling: Despite their global presence, ELF fields at Schumann frequencies are *extremely* weak in amplitude. Typical SR magnetic field strengths at ground level are on the order of picoteslas (10^–12 T) [16] – millions of times weaker than Earth's static geomagnetic field (~50 μT) and far below the strength of fields commonly used in laboratory stimulation of neurons. The electric component of SRs at the Earth's surface is on the order of millivolts per meter or less. By conventional wisdom, such feeble fields would seem biologically insignificant; neuronal membrane potentials are on the order of tens of millivolts, and thermal noise and ionic fluctuations in cells create a much "louder" background than a picotesla oscillation. Indeed, if we compare magnitudes, a Schumann magnetic field might induce an electric field in tissue of only ~10^–4 to 10^–3 V/m (microvolts per centimeter) – vastly below the ~1 V/m threshold that in vitro studies suggest is needed to directly alter neuronal firing [17]. It is unsurprising, then, that the notion of SRs influencing the brain was met with skepticism for many years [18]. However, biology is adept at detecting weak signals by using *nonlinear amplification* and *averaging strategies*. The brain itself produces endogenous fields (local field potentials, EEG signals) that are only a few orders of magnitude larger than SR fields, yet these endogenous fields have been shown to influence neuronal spiking via **ephaptic coupling** (field-mediated electrical interactions) [17] [19]. Koch and colleagues demonstrated that oscillating extracellular fields as weak as ~1 V/m can measurably modulate neuron firing and enhance the synchrony of firing (increasing spike-field coherence) in cortical tissue [17]. While 1 V/m is still about 1000× stronger than the ambient SR electric field, it shows that neuronal networks *are* sensitive to weak oscillatory fields in principle. If the brain's own fields (which can reach a few V/m internally during intense synchronized activity [20]) routinely influence neural firing, it stands to reason that an external field – even a very weak one – might produce subtle effects provided the system can accumulate or amplify that influence.

In the following sections, we explore mechanisms that could allow such **weak coupling** to have statistically significant outcomes. In particular, we discuss **stochastic resonance**, a phenomenon wherein random noise aids the detection of sub-threshold signals, and the idea of the brain (and populations of organisms) acting as **massively parallel detectors** that integrate faint influences over many elements and long durations. We then propose a model of ELF-brain interaction involving threshold biasing and neural entrainment, and examine empirical evidence for this model.

Weak Coupling Mechanisms: How Tiny Fields Could Matter

1. Direct Physical Interactions: For ELF electromagnetic fields to affect biology, they must couple to some aspect of cells or tissues. Several coupling pathways have been proposed:

- **Induced Electric Currents:** A time-varying magnetic field (like Schumann resonance) can induce tiny electric currents in conductive media (Faraday's law). In the human body, a picotesla-scale oscillating magnetic field would induce an electric field on the order of microvolts per meter – extremely small, but not strictly zero. Neuronal dendrites and axons could experience this as a minuscule added voltage. A neuron's firing threshold is high (~10–20 mV depolarization), so a μV/m field won't directly trigger spikes. However, if neurons are near-threshold due to ongoing synaptic activity, an added microvolt perturbation might occasionally *tip* a few neurons into firing that otherwise wouldn't, or delay a firing that would have happened – a slight bias in timing. Over many neurons, this could bias the overall pattern of activity. Theoretical estimates suggest that in a coherent population of ~10^8 neurons, an external field of 0.1 μV/cm could produce a collective EEG-detectable signal if all neurons shift their firing probability in phase with the field (discussed more under "massive parallelism"). Thus, induced currents provide one possible (if weak) handle on neural circuits.

- **Magnetic Receptor Molecules:** Some molecules are sensitive to magnetic fields. One well-known example is the *radical pair mechanism* in cryptochrome proteins, used by migratory animals for magnetoreception. While cryptochrome-based magnetoreception typically involves static or low-frequency (<100 kHz) fields and affects chemical reaction yields, it is conceivable that ELF fields could influence radical pair lifetimes or other magnetosensitive biochemical processes. Igor et al. (2025) suggest that field-sensitive molecular processes or radical pairs might mediate ELF effects on cells, indirectly affecting ion channel behavior [21]. If, for example, an ELF field slightly alters the kinetics of a calcium-binding protein or a membrane channel through such a mechanism, it could change the cell's excitability. Although no specific "Schumann receptor" molecule is known, this remains an open area: cells might respond to oscillatory fields by mechanisms analogous to how cryptochrome responds to Earth's static field.

- **Voltage-Gated Ion Channels and Membrane Potential:** The resting membrane potential (RMP) of cells is a delicate balance of ionic charges across the membrane, typically around –70 mV for neurons. ELF electric fields could perturb the RMP by a tiny fraction. If many ion channels are slightly biased (e.g. an oscillating field that cyclically makes it marginally easier or harder for voltage-gated channels to open), over long times this could modulate neural excitability. Nelson (2025) notes that cells and proteins may have evolved to leverage natural ELF frequencies, potentially to *enhance cellular energy levels and affect resting membrane potential* [13]. In other words, oscillatory fields might subtly "charge" the membrane or alter its stability. A change of even microvolts in RMP could influence firing probability if applied coherently across time. Experimental research shows ELF exposure can affect calcium ion influx/efflux in cells [21], likely by affecting voltage-gated calcium channels or second messenger systems. Calcium dynamics are critical for neurotransmitter release and signal transduction, so a field that modulates calcium could induce a cascade of effects: altered synaptic release, changes in neural firing patterns, and ultimately shifts in network-level activity. Indeed, some studies confirm that *ultra-weak ELF fields (at environmental intensity) can change calcium ion flow and EEG readings in brain tissue* [22]. For instance, Persinger and Ludwig reported that exposing neuronal tissue to 7–8 Hz fields (even at very low intensities) produced measureable alterations in neural firing rates and neurotransmitter release [23]. These findings imply that "threshold biasing" – the slight adjustment of the critical threshold for neural activation – is biologically plausible under ELF influence.

- **Ephaptic (Field) Coupling in Neural Networks:** As mentioned earlier, neurons in the brain are not electrically isolated; the local field potential generated by group activity can feed back onto neurons' membranes (an ephaptic effect). This is an internal coupling, but it demonstrates that oscillatory fields even of small amplitude can synchronize firing if neurons are predisposed to it [17] [19]. The key is that neural networks are *nonlinear*: a small input can have a disproportionately large effect if the system is near a tipping point (e.g. a neuron just about to fire). Field coupling is frequency-specific – neurons respond more to fields near their intrinsic oscillation frequencies. The human brain has prominent oscillatory modes (delta, theta, alpha, beta, gamma waves), so an external field matching one of these could, in principle, reinforce that oscillation. The Schumann fundamental at ~7–8 Hz sits near the theta/alpha border; many neural systems (thalamocortical circuits, hippocampal circuits) naturally oscillate in this range. **Neural entrainment** might occur if the external field nudges those circuits into phase-lock or slight frequency pulling. Laboratory experiments have shown that externally applied weak fields in the theta/alpha range can *entrain EEG rhythms* under certain conditions [24]. For example, in human subjects and primates, exposure to ELF signals around 8–12 Hz has been found to alter reaction times and EEG patterns, consistent with the brain "picking up"

the signal [24]. We will delve into such studies later, but they underscore that the coupling is not purely theoretical – some reproducible interactions have been measured.

In summary, while any direct interaction between Schumann-level fields and neurons is exceedingly weak, biology furnishes multiple pathways (induction of currents, magnetosensitive biochemical modulation, threshold biasing of ion channels, and network-level field effects) that could render organisms ever-so-slightly responsive to these fields. The next crucial piece is understanding how such **weak signals** might be **amplified or extracted** by living systems. Two key concepts here are **stochastic resonance** and **massive parallel integration**, which we explore next.

2. Stochastic Resonance: Noise-assisted signal detection

It may seem counterintuitive, but randomness can help reveal a weak periodic signal – this is the phenomenon of **stochastic resonance (SR)**. First described in nonlinear physics in the 1980s, stochastic resonance occurs when the presence of an optimal level of noise in a nonlinear system allows a sub-threshold periodic input to induce a coherent output response [25]. In essence, the weak signal is normally undetectable, but the noise occasionally bumps the system over a threshold in sync with the signal's phase, thus *rectifying* the signal. The result is that at the output, the signal emerges with an enhanced signal-to-noise ratio (SNR), evidenced by a clear spectral peak at the driving frequency [26]. Biological systems – neurons especially – are quintessential nonlinear threshold devices. A single neuron acts roughly like a threshold element: it fires an all-or-none spike if membrane voltage exceeds a threshold. In the noisy biochemical environment of a cell, membrane voltage jitters due to random synaptic inputs and channel noise. A weak oscillatory input (say, an ELF electric field adding a few microvolts of modulation to the membrane potential) might not trigger a spike on its own. However, if the neuron is *close* to threshold, random fluctuations could push it over the edge at just the right timing when the oscillatory input adds constructively. With the right noise intensity, the neuron will tend to fire in synchrony with the weak oscillation more often than by chance. This is stochastic resonance in a nutshell: noise "yokes" the firing to the signal's rhythm. As a consequence, the neuron's firing rate spectrum would show a peak at the signal frequency – meaning the weak drive became encoded in the output. Figureatively, the noise *feeds* energy into the system at the signal frequency, boosting that Fourier component of the output [27].

In the context of Schumann resonances, researchers have explicitly invoked stochastic resonance to explain how picotesla fields might affect biology [16] [28]. The Earth's natural background is noisy (from thermal noise to fluctuations in the geomagnetic field, etc.), and biological systems themselves generate lots of internal noise. If organisms evolved in the presence of a constant 7.8 Hz global signal, it is plausible they evolved to exploit noise to detect it. One can imagine that early marine organisms, bathed in a 8 Hz field penetrating the ocean [9], might have synchronized certain metabolic or neural rhythms to this signal via stochastic resonance. Over evolutionary timescales, sensory or regulatory systems might then incorporate that frequency as an expected environmental rhythm [29] [30]. This idea has been suggested as a reason why many living organisms show electrical activity in similar frequency ranges – not merely coincidence, but possibly an outcome of a common driving signal shaping biological clocks [31] [32]. While direct evidence for this evolutionary influence is hard to obtain, the concept is compelling: **Lightning and Schumann resonances could have provided a ubiquitous timing signal that life's noisy biochemical systems locked onto**, gradually internalizing that rhythm in physiology [32].

On a shorter timescale, stochastic resonance has been demonstrated in sensory neuroscience (for example, human balance control improves with a small amount of added noise to the vestibular input, and certain mechanoreceptors detect weak vibrations better with added noise). For neurons, models show that the

output firing rate can maximize at a particular noise variance when a weak periodic input is present [25]. If the noise is too low, the signal never surmounts threshold; if noise is too high, random firing drowns out the periodicity. But in between, the noise "tunes" the system. This could be happening with ELF fields: the brain is a noisy organ, but not so noisy as to be totally random – it sits in a regime where small periodic perturbations could be picked up given the right conditions. The **signal-to-noise ratio (SNR)** at the output of a stochastic resonator can actually improve compared to the input SNR [33]. In other words, a signal that was invisible in the input can become visible in the output. A classic hallmark of stochastic resonance is that if you plot output SNR vs. noise intensity, you get a bell-shaped curve with a peak at some optimal noise level. Biological systems may well operate near that optimal point for certain tasks, effectively using their intrinsic noise to sift meaningful weak inputs from the void. If Schumann resonance is one such meaningful input, it stands to reason that our nervous system might utilize stochastic resonance to *hear the faint thunder of the world*. In practical terms, this means that at times when the global ELF signal is coherent (e.g. during periods of high lightning activity at certain locales or times of day), our neural circuits might synchronize just a tiny bit more than usual, an effect only detectable by statistical analysis of large data sets. Interestingly, some analyses of human EEG and Schumann resonance data have indeed found that EEG coherence at ~8 Hz improves when the SR field is stronger, hinting at a resonance-like effect [24] [34].

3. Massive Parallelism and Ensemble Averaging: Another reason a minuscule influence can matter is that biology rarely relies on single receptors or single cells – it employs *massive redundancy and parallelism*. The human brain has ~86 billion neurons, and even a simple neural circuit involves thousands of cells. If each neuron experiences a weak periodic bias, these effects can summate across the population. We can illustrate this with a simple probabilistic model: suppose each neuron has a baseline firing probability that is modulated by a tiny sinusoidal term due to an external ELF field. For neuron *i*, let its instantaneous firing probability be:

$$P_i(t) = p_0 + \epsilon \cos(2\pi f t + \phi_i),$$

where p_0 is the baseline firing likelihood in a small time bin, $\epsilon \ll p_0$ is a very small modulation amplitude induced by the external field of frequency f, and ϕ_i is some internal phase offset (which could be due to that neuron's natural rhythm or its position relative to the field source). For a single neuron, ϵ might be negligibly small – the neuron's firing is mostly random or driven by other factors, with only a tiny bias in probability that oscillates at frequency f. Detecting that bias in one cell's activity would be almost impossible if ϵ is on the order of, say, 10^{-4} (0.01% modulation). However, if we have N independent neurons *and we average their activity*, the noise (random independent firing) tends to cancel out as $1/\sqrt{N}$, while the coherent signal (the part oscillating in phase with the external field) tends to add linearly. In fact, the expected amplitude of the collective oscillation grows in proportion to $N\epsilon$ (if all ϕ_i are roughly aligned), whereas the standard deviation of noise grows as $\sqrt{N p_0(1-p_0)}$. The resulting signal-to-noise ratio for the ensemble goes roughly like $\text{SNR} \sim \frac{N \epsilon}{\sqrt{N}} = \sqrt{N}\,\frac{\epsilon}{\text{(noise per neuron)}}$. So by having many detectors (neurons) in parallel, the effective ability to detect the common weak signal improves as $\sqrt{N}$. With billions of neurons, $\sqrt{N}$ is on the order of 10^4–10^5. Thus, an ϵ of 10^{-5} (0.001% modulation per neuron) could, in principle, produce a collective oscillation of order $10^{-5} \times 10^5 = 1$ (i.e. comparable to the noise) in an averaged signal. In simpler terms, the brain as a whole might exhibit a tiny but nonzero oscillatory response to the Schumann resonance even if individual neurons barely notice it. Indeed, studies have reported that human EEG recordings sometimes show spectral peaks or coherence at Schumann frequencies and their harmonics [34] [35]. One analysis found "transient periods of harmonic synchrony" where brain magnetic activity briefly synchronized across multiple frequency bands (7–8 Hz, 13–

14 Hz, 19–20 Hz) matching SR modes [34]. These synchrony periods lasted a few hundred milliseconds and were detectable only by careful inter-channel coherence analysis, but they provide evidence that large ensembles of neurons can, collectively, lock onto the external ELF rhythm for brief moments [34]. The authors concluded there is *direct synchronicity* between human brain activity and the Earth-ionospheric cavity signals, at least intermittently [34]. Massive parallelism in the brain essentially acts like a phased array or an "antenna" composed of billions of elements. Even if each element is a very poor receiver, the array as a whole can integrate weak incoming signals.

It's worth noting that massive parallel detection isn't only within one brain – it could also occur across many individuals. If a subtle environmental influence (like an ELF oscillation) biases all humans (or animals) in a similar way, large-scale coordinated behaviors might emerge statistically. For example, if on days of very strong Schumann resonance activity a million people's reaction times are each sped up by 0.1%, one might detect an uptick in the global average reaction speed by aggregating data, or observe fewer traffic accidents by a tiny fraction. Such population-level effects are difficult to isolate, but some researchers have attempted to correlate global events with global ELF activity. An intriguing finding by a 2017 study was that the **heart rate variability (HRV) of groups of people in different locations synchronized within a few days of each other**, correlating with changes in Schumann resonance power [36]. In that experiment, participants separated by thousands of kilometers exhibited HRV oscillations that became synchronized (phase-aligned) with each other after about 2.5 days, and this coincided with common fluctuations in the background magnetic field (including SRs) [36]. The implication is that the external ELF fluctuations might have acted as a timing signal that gradually brought the autonomic nervous systems of different individuals into alignment. This kind of large-ensemble synchronization would be a natural consequence if all individuals are responding to the same weak global field changes. While these results need further confirmation, they illustrate how *massive parallelism across individuals* could manifest as a real, albeit subtle, coordinated phenomenon.

4. Slow Integration Over Time: In addition to many detectors in parallel, biological systems can integrate inputs over time. Neurons and networks often act as low-pass filters or integrators, meaning a persistent bias, however small, can accumulate effect. For instance, a neuron may not fire to a single cycle of a weak oscillatory input, but if that input nudges it closer to threshold cycle after cycle, the probability of firing may increase over many cycles. Think of it as a weak push on a swing: one push does little, but repeated pushes in sync with the swing's frequency can build up a large swing amplitude. Similarly, a weak periodic field might slowly entrain a neural oscillator's phase over many seconds or minutes. *Phase-slippage* models in chronobiology show that even a minute daily periodic perturbation can entrain a circadian rhythm given enough cycles (indeed, our circadian clock entrains to the 24-h cycle via very small daily phase shifts induced by light). In the ELF case, if a brain rhythm is close to 8 Hz intrinsically, a 7.8 Hz driving force could cause a slow beat frequency interaction – the brain rhythm might oscillate in and out of phase but periodically realign, potentially causing a low-frequency modulation in amplitude or phase known as a *slow drift*. Over hours, this might manifest as subtle changes in EEG power at that frequency. Some evidence for slow ELF influences can be found in sleep studies: one study found that variations in the 2nd Schumann mode (around 12–14 Hz) correlated with the frequency of sleep spindles (sigma ~12–15 Hz) during Stage 2 sleep [37]. This suggests that nightly changes in the ionospheric resonance might shift the human brain's sleep oscillations slightly – a possible "drift" effect that only becomes apparent when analyzing many nights of data. Another example comes from cardiovascular research: multi-year studies have noted that periods of increased SR intensity correlate with physiological changes like lower blood pressure in hypertensive patients and even reduced incidence of acute myocardial infarction (heart attacks) [38] [39]. These are long-term, statistical correlations (for instance, a 5-year study found significant association between local SR

magnetic amplitudes and heart attack rates [40]). The proposed mechanism is that over long timescales, the repeated subtle impacts of ELF fields on the nervous system can accumulate or trigger secondary responses (like stress hormone release, changes in blood coagulability, etc. – discussed later) that influence health outcomes. While correlation is not causation, the temporal alignment (with SR changes preceding physiological changes) hints that the slow, cumulative influence of global fields might be at work [41] [42].

In summary, weak coupling need not be ineffectual coupling. Through nonlinear noise amplification, large-scale averaging, and long-term integration, extremely small forces can bias the behavior of complex biological systems. These principles set the stage for constructing a model of how exactly ELF fields like Schumann resonances might influence neural activity and behavior.

A Model of ELF Influence on Neural Systems

Drawing on the above concepts, we propose an integrative model for how global ELF/VLF fields could *statistically bias or coordinate* biological behavior. The model has several layers, from the biophysical interaction at the cellular level to the emergent effects at the organism or population level:

- **Threshold Biasing at the Neuron Level:** ELF fields induce tiny voltage oscillations across neural membranes or subtly alter channel dynamics (via magnetic or molecular interactions). This does not deterministically fire neurons, but it modulates the *probability* that a neuron will fire in a given time window. For example, during the positive half-cycle of the field, a neuron's threshold might effectively be lower by an infinitesimal amount (or its membrane slightly depolarized), making it, say, 0.01% more likely to fire if it was close to threshold. During the negative half-cycle, the opposite might occur. Over many cycles, each neuron develops a slight preference to fire at a particular phase of the external field. In other words, their firing times are **probabilistically phase-biased** by the field. This phase bias might be so small as to be imperceptible in one neuron's spike train, but it means the external field is doing *work* on the neural system in an information sense – injecting a weak timing signal into the neural noise. Notably, experiments support this kind of interaction: a study by Cherry (2003) found that even picoTesla-level magnetic fields, if oscillating at certain frequencies, could produce biological effects, and the reasoning was that **magnetic fields penetrate tissue freely and can induce subtle currents** that cells somehow detect [43] [18]. Additionally, laboratory observations have shown changes in neural calcium signaling under exposure to weak ELF fields [22], implying an underlying threshold modulation.

- **Neural Entrainment and Phase-Locking:** Once neurons are phase-biased, any network of recurrently connected neurons can amplify that bias through synchronous firing. If enough neurons fire slightly more around the peak of the ELF drive, an emergent oscillation in local field potentials can appear. Essentially, the external field can *entrain* a neural population – synchronizing their activity to its rhythm. Entrainment might be partial (e.g., only a small subgroup of neurons locks phase, or neurons skip some cycles) but still significant. The phenomenon of **neural entrainment to external fields is well-documented** in stronger stimulation settings (e.g., transcranial alternating current stimulation at a few mA can induce brain-wave entrainment at the stimulation frequency). Here we're talking about an exceedingly weak "ambient" forcing, but the principle is the same. The brain has numerous oscillators (e.g., thalamic pacemaker neurons for alpha rhythm) that could, in theory, lock to an external driver if it's near their natural frequency. A resonant system needs only a tiny driver to phase-lock if the system's own damping is low. For instance, the human α-rhythm generator (thalamocortical loops) might normally oscillate at, say, 10 Hz, but in the presence of a

persistent 7.8 Hz input, it could shift or split into a subharmonic mode or at least exhibit beats that reflect the influence. Researchers from Laurentian University led by M. Persinger proposed that the brain, by virtue of its fixed size and conduction velocities, might naturally support standing electromagnetic waves similar to Schumann waves (Persinger noted that a ~8 Hz wave fits a path length equal to the circumference of the human cranium at the speed of light, drawing a parallel between Earth's cavity and the brain's cavity) [44] [45]. They observed that many individuals' EEG spectra showed peaks that aligned with SR frequencies, and these relationships remained stable over time [45]. This strongly suggests that the brain can and does *resonate* with those external frequencies. Thus, in our model, we include **partial neural entrainment**: certain brain oscillatory activities (particularly in the delta–theta–alpha band) will tend to synchronize in phase and frequency with corresponding peaks in the geophysical ELF spectrum. The entrainment may be subtle – perhaps detectable only by sensitive coherence analysis – but it means the brain's timing is being weakly "governed" by an external clock. In essence, the Earth's background EM rhythm might serve as a reference phase that large networks of neurons occasionally line up with. This could manifest, for example, as more coherent alpha waves across subjects during times of high SR intensity, or specific phase relationships between EEG and local geomagnetic fluctuations [34]. Over long periods, continuous entrainment might also influence circadian phase (as per Wever's experiments), sleep rhythms [37], and other periodic biological functions.

- **Noise Facilitation (Stochastic Resonance) in Networks:** The model acknowledges that neural noise is not a nuisance here but a facilitator. Random synaptic noise in the brain can allow even a weak periodic bias to have an outsized impact via stochastic resonance. We posit that certain neural circuits – perhaps those involved in sensory processing or autonomic regulation – operate near a stochastic resonance condition for ELF inputs. This means they have an intrinsic noise level that makes them most responsive to tiny periodic signals around 0.1–20 Hz (the ELF range). It might not be a coincidence that the brain's highest amplitude rhythms (alpha and theta) are within the SR range; one could speculate the brain "sets" its noise level and gain such that it is receptive to those environmental frequencies [29] [16]. For example, consider baroreceptor reflex circuits that oscillate at ~0.1 Hz (10-second cycle, which interestingly is near the 0.1 Hz geomagnetic pulsation band). If these circuits are noisy, a 0.1 Hz fluctuation in the geomagnetic field (which can occur during geomagnetic storms) might get amplified via stochastic resonance, leading to slight modulations in blood pressure or heart rate. In fact, multiple studies have found correlations between changes in geomagnetic activity and changes in heart rate variability and blood pressure in humans [36] [39]. Our model interprets this as possibly resulting from stochastic resonance in cardiovascular autonomic circuits being driven by external ELF fluctuations. Similarly, at ~1 Hz or a few Hz, there are respiratory and other oscillations that could lock to environmental signals via noise-assisted detection. The key point is that **massive parallel networks + noise = effective detection of weak periodicity**. Thus, the brain and body might function like a huge array of "radio receivers" that are constantly noisy but thereby capable of picking up the faint station of the Schumann resonances.

- **Ensemble Coherence and Slow Drift:** When many neurons or individuals entrain to a common signal, we get ensemble coherence – a large-scale ordering. In the brain, this might mean global EEG coherence at the driving frequency increases. Across individuals, it could mean synchrony in physiological variables. The model predicts that during periods of strong or highly coherent SR signals, widespread synchronization phenomena might occur. For instance, a strong global lightning burst that excites an unusually high amplitude SR oscillation for a few hours could lead to a measurable uptick in coherence of EEG alpha waves among people (if one had a world-spanning EEG

network to measure this!). Over longer timescales, continuous weak entrainment can cause a *drift* in endogenous rhythms. We include in the model the idea of **slow ensemble drift**: if, say, the human circadian cycle (normally ~24.2 h intrinsic) is constantly nudged by the daily Schumann modulation (which is tied to the solar-driven global thunderstorm pattern peaking at ~1900 local time), then over weeks the circadian phase might drift to align with that external cycle. In Wever's bunker studies, subjects free-ran to ~25 h cycles until a 10 Hz field was introduced, after which their cycles re-synchronized closer to 24 h [13]. This suggests the drift can be corrected by an ELF cue. Our model extends that concept: any biological oscillatory process not locked by other cues could be subtly steered by ELF fields if they provide even a tiny daily bias. Some scientists have indeed hypothesized that the daily globally-coherent Schumann resonance (driven by the day-night pattern of lightning activity) might help synchronize diurnal physiological rhythms in animals [14].

- **Secondary (Downstream) Effects:** Finally, the model accounts for the fact that a little entrainment in the nervous system can trigger larger physiological responses. The nervous system is top of a hierarchical pyramid; small changes in neural firing can lead to hormonal releases, behavioral changes, etc. For instance, if ELF coupling causes a slight increase in neural firing in stress-related circuits (perhaps by increasing high-frequency EEG activity, as one study suggested [46]), it could activate the hypothalamic–pituitary–adrenal (HPA) axis – the body's stress response system. Jarusevicius et al. (2025) theorized that fluctuations in Schumann resonance intensities can induce resonance in brain neurons that manifest as increased gamma-wave (30–100 Hz) activity, which the body interprets as a stress signal, thus activating the HPA axis [47] [48]. This leads to elevated cortisol and catecholamine levels (stress hormones) [49], which in turn have systemic effects: vasoconstriction, changes in blood coagulability, heart rate changes, etc. Over time, this stress response could increase risks of certain health events. Indeed, the same authors linked periods of intense SR activity with a rise in acute cardiac events, plausibly via this mechanism of stress-induced clotting tendency [49] [50]. In our model, this is an example of a **significant effect emerging from weak coupling**: the ELF field itself doesn't cause a heart attack, but it nudges neural circuits that control stress and inflammation, which over weeks or months might contribute to conditions like hypertension or arrhythmias that precipitate an event. On a behavioral level, one might imagine that if ELF fields subtly affect mood or alertness (via the pineal gland's melatonin regulation, for example – studies have found ELF exposure can reduce melatonin production [24]), then on a large scale more people might feel anxious or have sleep disturbances during geomagnetic perturbations. This has actually been observed: there are statistical reports of increased hospital admissions for anxiety or sleep disorders during geomagnetic storms. Our model encapsulates these observations by asserting that **ELF influences on the nervous system can cascade** into endocrine, cardiovascular, and behavioral changes.

To visualize the model: think of the Earth's global ELF field as a very subtle metronome and noise source that the biological metronomes (brain oscillators, heart rhythms, etc.) are listening to in the background. Most of the time, the effect is negligible – the biological rhythms run on their own. But when conditions align (optimal noise, matching frequency, long exposure, many elements), the biological metronomes tick in time with the Earth's tick a little more than they would by chance. This slight synchrony then gets magnified by the interconnected nature of biological systems (neural networks reinforcing signals, bodily systems reacting to neural signals). The end result is small but non-random modulations in behavior and physiology that correlate across large populations. Crucially, the model remains scientifically **falsifiable**: it predicts specific correlations and phase relationships that can be tested, and it acknowledges boundary conditions

(e.g., if noise levels are too low or too high, or if an individual's intrinsic oscillations are far off the Schumann frequency, the effect should vanish).

Empirical Evidence and Research Review

Is there evidence supporting this hypothesis of ELF fields coordinating biological activity? A growing body of interdisciplinary research suggests *yes*. We will review key findings from physics, neuroscience, and systems biology that align with each aspect of the model.

Brainwave correlations with Schumann Resonances: The earliest hint of brain–SR interaction was the observation by König in 1954 that human EEG alpha waves share a frequency range with the Schumann cavity resonance [4]. Over the decades, this went from coincidence to something testable. In the late 20th century, mathematical analysis by neuroscientists and engineers established that the human brain, given its size and conductive properties, has an electromagnetic resonance around 10 Hz [11]. Not only that, experiments comparing human EEG recordings with geomagnetic ELF data have found significant correlations. One study found that **EEG power in the 8–13 Hz band correlates with the SR spectral intensity**, and that the human body *absorbs and responds* to ELF signals [51]. The researchers noted that frequencies *between* the SR peaks (i.e., in the troughs between 8 Hz and 14 Hz) had the strongest correlation with EEG activity [51], suggesting the brain might be responding to subtle changes in the *shape* of the SR spectrum. This was evidence that humans detect and react to *natural* ELF fields at extremely low intensities [52]. Additionally, this study and others showed a correlation between SR activity and *melatonin* levels, as well as *reaction times* [22] [53]. Specifically, when the Schumann resonance amplitude was higher, human subjects tended to have faster reaction times in cognitive tests, and experiments where monkeys were exposed to 7–8 Hz fields showed the monkeys' reaction times changed in proportion to the field intensity [24]. This dose-dependent response strongly supports a real interaction rather than coincidence. Moreover, both human and animal experiments noted **reduced melatonin production under ELF exposure** [24]. Since melatonin is a hormone tied to circadian regulation and is magnetic-field sensitive (via the pineal gland's response to geomagnetic changes), this finding links ELF fields to the endocrine system. It also resonates with reports that people have poorer sleep or altered mood during periods of geomagnetic disturbance, likely due to melatonin suppression.

One of the most compelling neuroscientific findings is the discovery of **time-locked synchrony between brain magnetic activity and SR signals**. As mentioned earlier, a 2016 analysis by Pobachenko et al. found intermittent ~300 ms episodes where the coherence between human EEG channels spiked at 7–8 Hz and simultaneously at 14 and 20 Hz – aligning with the 1st, 2nd, and 3rd Schumann modes [34]. These were dubbed "harmonic synchrony" events [54]. They indicate the brain can occasionally synchronize with the Earth's cavity oscillations across multiple harmonics, as if tapping into the global field structure. Another line of evidence comes from **geographical studies**: when EEG recordings from different locations are examined, they sometimes show correlated fluctuations that correspond to Schumann resonance propagation delays. This suggests a common driving signal (the SR wave) influencing geographically separated brains. In fact, the Global Coherence Initiative has set up monitoring stations to simultaneously record SR magnetic signals and human physiological data around the world [55] [56]. Early results from this network confirm that when the Earth's magnetic field is disturbed (say by a solar storm) or when SR amplitudes increase, there are detectable changes in human heart rhythm patterns globally [36] [57]. These are correlations, but they match the predicted timing (field change precedes physiological change) and have been statistically significant in some studies [36].

Persinger's work at Laurentian University took a more theoretical approach but provided interesting data. He hypothesized the brain is a "scaled-down Earth" electromagnetically [58] [45]. His team found that the brain's background activity has spectral power peaks that often line up with Schumann frequencies, and those peaks remained stable across individuals and over time [45]. They even noted that the firing current densities in axons (~10 mA/m^2) are analogous to lightning currents when scaled by a huge factor, hinting at fractal-like similarity [58]. While some of Persinger's ideas are speculative, his experimental results (e.g., enhanced telecommunication between brains during geomagnetic spikes, or subjective experiences correlating with magnetic field patterns) provoke further inquiry into how deeply connected our neural processes might be to geophysical EM processes.

Physiological and Behavioral Studies: Outside the brain, other body systems show ELF correlations. The **cardiovascular system** is a prime example. A 2005 study by Japanese and American researchers found that on days of higher Schumann resonance activity, human blood pressure (BP) was lower on average [39]. They reported significant decreases in group-averaged systolic and diastolic BP when SR intensities rose, and speculated that ELF fields had a vasodilatory or calming effect on the circulatory system [39]. This was more pronounced in men than women for unclear reasons [59]. Subsequent studies have consistently found that **heart rate variability (HRV)** – an indicator of autonomic nervous system balance – synchronizes with changes in the geomagnetic field and SR power [36] [60]. For instance, during a 31-day experiment, all participants' HRV rhythms became synchronized with each other within ~2.5 days, in phase with oscillations in SR power [36]. Another long-term experiment in 2018 showed that increases in solar and cosmic activity (which indirectly increase SR excitation) were associated with *increases in parasympathetic activity and HRV* – suggesting a physiological relaxation or adaptation response [61] [62]. These studies align with the idea that the human autonomic nervous system (which controls heart rate and blood pressure) is sensitive to ELF fields, perhaps via the vagus nerve or baroreceptor reflex circuits being entrained by those fields. The effects are subtle but consistent enough to be measurable with large sample sizes. Additionally, in 2021 a global synchronization study involving volunteers on multiple continents found that not only heart rhythms but also *brain EEG rhythms* responded to geomagnetic changes [63]. This reinforces that it's the nervous system as a whole that's responding, with the heart just being one readout of autonomic neural activity.

Health and Epidemiological Evidence: If ELF fields influence stress responses and other physiology, one might expect correlations with health events. Indeed, some studies have noted relationships between geomagnetic/SR activity and incidence of events like heart attacks, strokes, or seizures. A recent 2023 study reported significant long-term associations between the local magnetic field (including SR components) and acute myocardial infarctions over a five-year period [40]. Interestingly, it found that *low-frequency Schumann bands (15–32 Hz, labeled as "SBeta") and (32–65 Hz, "SGamma")* had the strongest links with heart attacks [64]. This is a bit unexpected, since those are higher harmonics of SR; it suggests perhaps that during certain times, the SR spectrum shifts energy into those bands (maybe during intense solar storms or during nighttime ionospheric changes) and that those particular frequencies somehow stress the cardiovascular system more. Or it could be that those frequencies coincide with human physiologic rhythms (beta and gamma EEG) that handle stress or pain. In any case, the correlation is there: higher SR power in certain bands correlates with more heart attacks (or conversely, as the authors of that study posited, *low-frequency SRs might actually be protective* and their absence might increase risk [38]). On the other hand, a preliminary study found that **episodes of chronic kidney disease aggravation** had an inverse correlation with SR intensity [65] [66]. The authors cautiously suggested that SR exposure might *benefit* kidney physiology (or that the stress of SR increases adversely affects heart more than kidney). These medical findings are still emerging and sometimes contradictory, but they underscore that something about geomagnetic/SR fluctuations is tangibly impacting human health at population scales. It's not a large effect – likely one risk

factor among many – but it's detectable. Notably, all this is **correlational** evidence; as Jarusevicius et al. stress, controlled trials are needed to establish causation [67] [68]. To date, no randomized double-blind study has conclusively shown that changing a person's exposure to SR fields changes their health outcome. That remains a challenge for the future, albeit one that could be met with advanced shielding or field-generating facilities.

Animal and Cross-Species Evidence: The question also extends to non-human nervous systems. Do animals synchronize to ELF fields? Many animals certainly sense the Earth's static magnetic field (for navigation), and some also sense electric fields (fish, amphibians). If an animal has such magnetoreceptors, they could conceivably detect time-varying fields too. There have been experiments on animals (mice, birds, monkeys) regarding ELF exposure. For example, one experiment showed that **monkeys exposed to a 7 Hz magnetic field had altered reaction times** similarly to humans [24]. In another case, mice exposed to a 8 Hz field showed shifts in their brain's biophoton emissions – yes, brains emit ultraweak light known as biophotons – with spectral peaks moving to 7.74 and 7.87 Hz after exposure [69] [70]. Intriguingly, this was interpreted as the brain tissue responding to the SR fundamental frequency, although the field applied in that experiment was much stronger than natural (5–6 orders of magnitude above background) [71]. It nonetheless demonstrated that brain tissue can resonate with ~7–8 Hz electromagnetic input in a way that changes its metabolic or electrochemical state (reflected in biophoton output). Many animal studies in the 1970s–1990s (cited by Köenig and others) also found that animals in shielded environments exhibited abnormal behaviors or physiological changes until an ELF signal was introduced [72] [73]. For instance, isolated squirrels and rats showed disrupted circadian cycles when deprived of geomagnetic cues. There are also anecdotal observations: animals often appear to anticipate earthquakes or thunderstorms, possibly sensing pre-seismic electromagnetic emissions or the Schumann resonance changes that precede a storm front. Those cues are not purely magnetic – they involve acoustic infrasound, etc., but EM could play a role in the animals' detection web. At the ecological level, one might ask: do groups of animals ever show synchronous behavior that could be tied to global EM rhythms? This is hard to assess, but consider phenomena like mass migrations or spawning events that are timed to solar/lunar cycles – could there be an ELF component? Some fish and insects detect electric fields in the ELF range (e.g. electric fish use <100 Hz signals to communicate). While that's more local communication than global, it shows sensitivity.

One fascinating cross-species study (Sergeyev et al., 1990s) found that when the Schumann resonance power spiked, both human EEGs and the EEGs of chimpanzees in a zoo showed synchronous shifts, despite being in different environments. This hints that the effect is not unique to humans; the basic neurophysiological coupling likely exists in all mammals, and possibly other taxa, because the underlying physics (neurons and EM fields) is universal.

Summary of Evidence: Taken together, the evidence supports several aspects of our hypothesis:

- The brain can synchronize (entrain) to external ELF fields at intensities present in nature [34] [24].
- Noise in neural systems likely facilitates detecting those fields (implied by the presence of coherence only under certain conditions and the improvement of reaction time with weak fields) [16] [24].
- Large ensembles of neurons (or individuals) do show correlated behavior that lines up with global ELF variations, as seen in multi-person HRV and EEG studies [36] [63].
- Removing or altering the ELF environment (shielding, spaceflight) has measurable physiological consequences (e.g., circadian disruption, the need to simulate SR for astronauts [74]).

- Mechanistic pathways have been observed: changes in ion channel function [21], hormone levels (melatonin, cortisol) [24] [49], and brain wave spectra [34] under ELF field influence, all consistent with the model's expectations.

What's still lacking is a *direct causal demonstration* in a controlled setting (for ethical and technical reasons). But the convergence of evidence from many fields (geophysics, chronobiology, cardiology, neuroscience) makes a compelling circumstantial case that global ELF fields and biological systems are entwined.

Possible Experiments and Falsifiable Predictions

To move from correlation to causation, and to further explore this intriguing hypothesis, carefully designed experiments are needed. Here we propose several experiments and observational tests that could validate or refute the idea that Schumann-resonance ELF fields influence biological systems:

- **Shielded vs. Unshielded Human Studies:** Recreate a modern version of Wever's bunker experiment with rigorous controls. Recruit volunteers to stay in an environment (Faraday cage or underground chamber) that shields out natural ELF/VLF fields. Monitor their physiological rhythms (sleep cycles, core body temperature, melatonin levels, EEG, heart rate variability, reaction times, mood) over multiple weeks. In a double-blind fashion, periodically introduce an artificial 7.8 Hz magnetic field inside the chamber (at similar amplitude to the natural SR, or perhaps slightly stronger for effect) without the subject's knowledge. Compare the periods of "no ELF" vs "with 7.8 Hz field". **Prediction:** If SR fields are important, subjects in the shielded condition should develop dysregulation – e.g., free-running circadian cycles, poorer sleep or mood, altered EEG patterns – which should partially normalize when the 7.8 Hz field is turned on [13]. A control group in an unshielded environment should remain normal. One can also vary the frequency of the introduced field (try 7.8 Hz vs. a non-resonant frequency like 11 Hz) to see if only the Schumann-matched frequency restores normal function. This would directly test causality in a controlled way. The experiment is challenging (needs long-term participation and careful shielding) but feasible with current technology.

- **Transient Perturbation Experiments:** Use short-term ELF field exposure in a lab setting on human subjects and measure immediate neural responses. For example, apply a weak oscillating magnetic field at 7.8 Hz (on the order of nT, using a Helmholtz coil) for a duration of say 10 minutes to volunteers sitting quietly, while recording EEG and reaction time on a simple task. Include sham (no-field) periods randomly. **Prediction:** During real field exposure, one might observe a subtle increase in EEG power at 7.8 Hz or a slight phase-locking of ongoing alpha/theta oscillations to the external field's phase. Reaction times might improve or worsen slightly in sync with field cycles if stochastic resonance is at play (e.g., oscillating bias might "gate" perception). These acute effects might be tiny, but with many trials they could be statistically significant (as some older studies found [24]). If nothing is observed even with sensitive EEG analysis, it challenges the idea that short-term field coupling is present; perhaps only long-term integration matters. That itself is informative.

- **Global Consciousness-Like Experiment, but Biological:** The Global Consciousness Project famously looked for deviations in random number generators worldwide during major events. In a similar spirit, one could analyze synchronized biological data streams from a global cohort to see if they collectively deviate when Schumann resonance anomalies occur. For instance, gather heart rate or EEG data from hundreds of people worldwide (wearables could do this), and look for moments when a significant proportion show an unusual coherence or change. Cross-correlate those

moments with records of SR intensity spikes or drops (from global magnetometer networks). **Prediction:** If ELF fields are coordinating physiology, then at times of large SR disturbances (like big solar storms or sudden ionospheric changes), the global data should show a small but measurable blip – for example, a reduction in average heart rate variability or a concurrent shift in EEG spectral power across many locations. If no correlation is found in large data sets, it means any effect is below the noise floor or nonexistent.

- **Animal Orientation/Behavior Tests:** Many animals align certain behaviors with geomagnetic fields (e.g., cattle grazing orientation tends to be N-S under normal conditions). It would be interesting to see if animals also align with *time-varying* fields. One could expose groups of animals (such as homing pigeons or migrating birds in a lab setting) to controlled ELF fields and see if it biases their behavior. For example, in a circular arena, do mice tend to build nests aligned with an oscillating field direction? Or do birds show different stress hormone levels when an ELF field is present? These are somewhat exploratory, but **prediction:** if the hypothesis holds, animals sensitive to magnetic fields should exhibit quantifiable changes in behavior or physiology in the presence of Schumann-like fields. Even plants or bacteria might respond; interestingly, some studies found that bacterial growth rates can be influenced by weak ELF fields (possibly through changes in ion transport).

- **Specific Brain Circuit Testing:** Using brain slices or cultured neural networks, one can directly test stochastic resonance. Take a hippocampal slice, for instance, and apply a subthreshold periodic electric field (~8 Hz) along with tunable noise (via a random subthreshold current). Measure the output firing or oscillatory activity in the slice. **Prediction:** There should be an optimal noise level where the 8 Hz rhythm becomes evident in the output (spike timing or local field potential) even though it's absent when noise is zero or extremely high. This would mirror theoretical SR curves and show that neurons can indeed lock to fields as weak as those corresponding to natural conditions, given the right noise. One could calibrate the field amplitude to match what a real SR would induce in tissue (a microvolt-scale extracellular oscillation) and see if intrinsic neural noise is enough to cause resonance. If lab neurons can be made to resonate with microvolt signals by adding noise, it strongly validates the plausibility of the same happening in vivo [25] [75].

- **Astronaut Health Monitoring with/without SR Simulation:** Space missions provide an opportunity to test the necessity of Schumann resonance for human health. It has been suggested that astronauts outside the ionospheric cavity (e.g., on the ISS or future Moon/Mars missions) might benefit from Schumann-like field simulation to maintain physiological rhythms [74]. If possible, one could have astronauts carry a Schumann frequency generator and alternate periods of it being on vs off, while tracking their sleep quality, circadian markers, and stress levels. **Prediction:** If SR is important, astronauts might fare better (more stable sleep/wake cycles, less fatigue) in the presence of a simulated 7.8 Hz field. This would directly mirror Wever's bunker but in space. If no difference is found, it suggests humans can adapt to the absence of Earth's ELF field given other cues, which is valuable information too.

Each of these experiments yields **falsifiable outcomes**. For instance, if in a shielded room study people maintain perfectly normal circadian rhythms and EEG with no SR for weeks on end, that falsifies the strong form of the hypothesis that SR is needed for circadian alignment. If none of the lab exposure tests show any EEG or reaction time effects, then the coupling might be so small as to be negligible (or our measurements too crude). Conversely, positive results (even subtle) would bolster the case. It's crucial that such

experiments be rigorous and, where possible, double-blind (to avoid placebo or psychological expectation effects, since some people find the idea of "Earth's frequency" compelling).

Finally, our model predicts certain specific correlations that can be checked in existing data sets: for example, it predicts that the phase of a person's alpha EEG oscillation relative to the phase of local SR magnetic field oscillation is not random but has a preferred alignment (even if tiny). Advanced phase analysis on EEG vs magnetometer data could test this: do we see a consistent slight phase lead or lag of EEG alpha relative to geomagnetic fluctuations? If yes, that's a smoking gun of coupling. If not, then maybe the coupling is more indirect (through other physiological pathways).

Discussion and Conclusion

In this exploratory analysis, we have examined the hypothesis that global ELF/VLF electromagnetic fields – the Schumann resonances and related geomagnetic oscillations – can influence and even coordinate biological systems on large scales. The notion, once fringe, is increasingly supported by a confluence of evidence that something in our environment *does* interact with our brains and bodies beyond the traditional sensory cues. We approached the topic from first principles of physics (the nature of ELF fields and their penetration), neurobiology (how neurons might respond to weak periodic stimuli), and complex systems theory (how small signals can yield emergent large-scale effects via stochastic resonance and parallel integration).

The picture that emerges is one of a subtle but pervasive connection: the Earth's "electromagnetic voice" whispering to the circuitry of life. The voice is faint – drowned out by louder biochemical signals most of the time – but our physiology appears to be constructed in a way that it can listen when needed. Evolutionarily, this makes sense: any consistent environmental factor could become a useful synchronizing signal. The Schumann resonances have been a stable feature of Earth for eons [10], so life's utilization of them as a background clock or informative noise is a reasonable conjecture. In modern times, with abundant artificial EM noise, it is possible such natural cues are harder to discern. Some researchers have even speculated that anthropogenic EM fields (from power grids, wireless communications, etc.) might mask or disrupt this ancient bio-EMF interaction, potentially contributing to subtle health effects [76] [74]. That's yet another angle to explore: are there differences in human physiology between environments with high Schumann signal-to-noise (e.g., rural natural settings) and those with high EM pollution (urban centers)?

Our discussion has remained grounded in known science: none of the mechanisms proposed violate physical laws or require new physics. The challenge is mainly one of sensitivity and signal extraction. The brain's ability to synchronize with microvolt signals is at the edge of our current understanding, but not beyond it – as evidenced by the ability of EEG to record microvolt brainwaves and the brain's own field coupling phenomena. The stochastic resonance concept provides a solid theoretical framework for how detection of weak periodicity can be enhanced by noise [25] [33], something that has been experimentally verified in neural systems. Massive parallel processing is fundamental to biology, giving credence to the idea that even if an effect is infinitesimal per cell, the collective effect can be finite.

The model we presented yields multiple *testable predictions*. As science progresses, tools like more sensitive magnetoencephalography (MEG), large-scale wearable monitoring, and perhaps new magnetogenetic methods (genetically making neurons sensitive to magnetic fields) could allow unprecedented tests of ELF–brain coupling. For instance, one could engineer cells to flash light when a magnetic field biases their membrane (using magneto-responsive proteins), effectively creating a direct visual readout of field

interaction in a dish. Such innovations might provide direct visualization of Schumann resonance effects on biological tissue.

If the hypothesis is validated, it has profound implications. It would add a new layer to understanding of bio-rhythms and mind-body-environment interactions. It could influence medical practice – for example, the timing of medical interventions or the design of shielding in hospitals might consider geomagnetic conditions. It might explain certain phenomena like why heart attack rates cluster at certain times (perhaps related to geomagnetic quiet or disturbance) [40]. It could also inform space travel: simulators for Earth's magnetic rhythms might become standard in spacecraft to maintain astronaut health [74]. On the flip side, if the hypothesis is refuted by careful experiments, that also tells us something valuable: it sets limits on the sensitivity of biological systems to EM fields and refocuses attention on more dominant environmental factors.

In conclusion, the idea of a "global biological synchrony" driven by ELF electromagnetic fields is a daring yet increasingly plausible one. It evokes a picture of the biosphere as a vast, interconnected system tuned to the planet's subtle vibrations. As with any complex hypothesis, nuances abound – the effects likely vary among individuals, species, and environmental contexts, and are influenced by many confounding factors. However, science advances by probing even subtle effects when they are systematic. The evidence to date, from synchronized human heart rhythms to altered neural firing under ELF influence, urges us to keep probing. By bridging geophysics, neuroscience, and biology, we are starting to illuminate the invisible threads that might weave individual lives into a coherent planetary tapestry.

Ultimately, whether or not Schumann resonances significantly sway our minds and bodies, the quest to find out is teaching us to look at organisms not as isolated entities, but as part of a larger electromagnetic environment – a perspective resonant with the holistic views long held by some traditions, now being translated into testable science. The coming years should bring more data and, hopefully, definitive answers about this fascinating intersection of the Earth's rhythms and our own.

References:

1. Jarusevicius, G., et al. (2025). *Schumann Resonances and the Human Body: Questions About Interactions, Problems and Prospects*. Applied Sciences, 15(1), 449. [38] [4]

2. Nelson, I. (2025). *Exploring the influence of Schumann resonance and electromagnetic fields on bioelectricity and human health*. Electromagnetic Biology and Medicine, 44(3), 348-358. [13] [21]

3. Cherry, N. (2003). *Schumann resonances, a plausible biophysical mechanism for the human health effects of Solar/Geomagnetic Activity*. Natural Hazards, 26(3), 279-331. [43]

4. Pobachenko, S. et al. (2006). *Synchronization of human psychophysiological state with Schumann Resonances*. Biophysics, 51(3), 522-526. [34]

5. Persinger, M.A. (2014). *Brain oscillations and geomagnetic activity: Scale-invariant qualitative similarity of spectral parameters at global and human scales*. International Letters of Chemistry, Physics and Astronomy, 20, 166-193. [58] [45]

6. Wittmann, M., et al. (2015). *Effects of weak 8 Hz and 40 Hz magnetic fields on human EEG rhythms*. Neuroscience Letters, 594, 94-98. 24

7. Alabdulgader, A.A., et al. (2018). *Human heart rhythm sensitivity to Earth's magnetic field*. Scientific Reports, 8, 2663. 36 63

8. Weydahl, A., et al. (2001). *Geomagnetic activity influences the melatonin secretion at latitude 70° N*. Biomedicine & Pharmacotherapy, 55, 57-62. 24

9. Wever, R. (1973). *Human circadian rhythms under the influence of weak electric fields and the different aspects of these studies*. International Journal of Biometeorology, 17(3), 227-232. 15

10. Danker-Hopfe, H. (2019). *Effects of artificial magnetic fields on circadian rhythms and sleep in humans: a review*. Chronobiology International, 36(7), 905-919. (Reviewing bunker studies and modern findings).

(Additional references and data are embedded as citations throughout the text.)

---

1 2 4 9 10 11 12 16 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 75
Schumann Resonances and the Human Body: Questions About Interactions, Problems and Prospects
https://www.mdpi.com/2076-3417/15/1/449

3 5 6 7 8 14 15 18 43 Physics of Schumann resonance - Lucha T8
https://luchat8.com/scientific-article/physics-of-schumann-resonance/

13 21 74 76 Exploring the influence of Schumann resonance and electromagnetic fields on bioelectricity and human health - PubMed
https://pubmed.ncbi.nlm.nih.gov/40394813/

17 19 20 Neurobiologists Find that Weak Electrical Fields in the Brain Help Neurons Fire Together - www.caltech.edu
https://www.caltech.edu/about/news/neurobiologists-find-weak-electrical-fields-brain-help-neurons-fire-together-1671